

# Analyzing biases of nitrogen contents and $\delta^{15}\text{N}$ values arising from acidified marine sediments with different $\text{CaCO}_3$ concentrations

PENG Yajun<sup>1, 2</sup>, LIU Dongyan<sup>1, 3\*</sup>, WANG Yujue<sup>1</sup>, RICHARD Pierre<sup>4</sup>, KEESING John K.<sup>5</sup>

<sup>1</sup> Key Laboratory of Coastal Zone Environmental Processes and Ecological Remediation, Yantai Institute of Coastal Zone Research, Chinese Academy of Sciences, Yantai 264003, China

<sup>2</sup> School of Management Science, Guizhou University of Finance and Economics, Guiyang 550000, China

<sup>3</sup> State Key Laboratory of Estuarine and Coastal Research, East China Normal University, Shanghai 200062, China

<sup>4</sup> Littoral, Environment et Sociétés, UMR 7266 CNRS-Université de La Rochelle, 2 rue Olympe de Gouges, La Rochelle 17000, France

<sup>5</sup> CSIRO Oceans and Atmosphere and University of Western Australia Oceans Institute, Indian Ocean Marine Research Centre, Crawley 6009, Australia

Received 26 May 2017; accepted 30 August 2017

© Chinese Society for Oceanography and Springer-Verlag GmbH Germany, part of Springer Nature 2018

## Abstract

In some studies, the researchers pretreated and measured organic carbon, nitrogen and their isotopes ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) of marine sediment together, to save costs and resources of analysis. However, the procedure of acidification to remove inorganic carbon for analysing  $\delta^{13}\text{C}$  can affect the values of nitrogen and  $\delta^{15}\text{N}$ , and the biases vary a lot depending on the  $\text{CaCO}_3$  contents of sediments. In this study, the biases of total nitrogen (TN) and  $\delta^{15}\text{N}$  values arising from acidified sediments were compared between the  $\text{CaCO}_3$ -poor (1%–16%) and  $\text{CaCO}_3$ -rich (20%–40%) samples. TN and  $\delta^{15}\text{N}$  values were altered during acid treatment (without centrifugation) that possibly led to N-containing compounds volatilization. For  $\text{CaCO}_3$ -poor samples, acidification led to a range of 0%–40% TN losses and 0‰–2‰ shift in  $\delta^{15}\text{N}$  values; and 10%–60% TN losses and 1‰–14‰ shift in  $\delta^{15}\text{N}$  values for  $\text{CaCO}_3$ -rich samples. The biases from most samples exceeded the precision of the instrument (0.002% for TN and 0.08‰ for  $\delta^{15}\text{N}$ ), and high biases could mislead our judgment for the environmental implication of the data. Thus, avoiding co-analysis of organic carbon, nitrogen and their isotopes ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) in sediments, even for  $\text{CaCO}_3$ -poor marine sediments, is necessary.

**Key words:** stable isotope nitrogen, nitrogen,  $\text{CaCO}_3$ , acidification, marine sediment

**Citation:** Peng Yajun, Liu Dongyan, Wang Yujue, Richard Pierre, Keesing John K. 2018. Analyzing biases of nitrogen contents and  $\delta^{15}\text{N}$  values arising from acidified marine sediments with different  $\text{CaCO}_3$  concentrations. *Acta Oceanologica Sinica*, 37(8): 1–5, doi: 10.1007/s13131-018-1188-2

## 1 Introduction

Organic matter contents (e.g., total carbon and nitrogen) and their isotope fingerprints ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) in the sediments are important environmental proxies to evaluate the sediment quality and trace the material sources in coastal and estuarine ecosystems (Tucker et al., 1999; Voss et al., 2000; Savage, 2005). For example, anthropogenically enriched organic nitrogen (e.g., human sewage, livestock runoff) can significantly elevate the values of  $\delta^{15}\text{N}$  to 10‰–25‰ (Sweeney and Kaplan, 1980; Savage, 2005). In some studies, the researchers pretreated and measured organic carbon, nitrogen and their isotopes ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) in marine sediment together, to save costs and resources of analysis (e.g., Middelburg and Nieuwenhuize, 1998; Kennedy et al., 2005; Hu et al., 2006; Wang et al., 2013). Acidification is a necessary procedure to remove inorganic carbon for meeting the requirement of total organic carbon (TOC) and  $\delta^{13}\text{C}$  measurements. However, when samples were acidified, the loss of total nitrogen (TN) and

even the shift of  $\delta^{15}\text{N}$  values occurred (Bunn et al., 1995; Jacob et al., 2005; Jaschinski et al., 2008; Serrano et al., 2008; Schlacher and Connolly, 2014). This biases caused by acidification varied greatly depending on the species of acid reagent (e.g., HCl,  $\text{H}_2\text{SO}_3$ ,  $\text{H}_3\text{PO}_4$  or HF) and carbonate contents in the sediments (Schubert and Nielsen, 2000; Lorrain et al., 2003; Kennedy et al., 2005; Schmidt and Gleixner, 2005; Fernandes and Krull, 2008). For examples, Harris et al. (2001) only detected a slight increase in  $\delta^{15}\text{N}$  values (0.04‰–0.14‰) after acidification using HCl fumigation; Fernandes and Krull (2008) also found 1‰–2‰ deviation of  $\delta^{15}\text{N}$  values between the acidified (using HCl washing) and non-acidified samples. For the biases of nitrogen content, Lohse et al. (2000) reported there were approximately 28%–47% losses in nitrogen compounds for  $\text{CaCO}_3$ -rich (55%–61%) samples after acidification using  $\text{H}_2\text{SO}_3$ ; Kennedy et al. (2005) however only found about 0.3% nitrogen losses in the acidified samples (30%  $\text{CaCO}_3$ ) using HCl. Therefore, there is uncertainty

Foundation item: The National Natural Science Foundation of China under contract No. 41376121; the Strategic Priority Research Program of the Chinese Academy of Sciences under contract No. XDA11020405; the Natural Science Foundation of Shandong Province under contract No. JQ201414; the Program of Advanced Talents of The Guizhou University of Finance and Economics, 2016.

\*Corresponding author, E-mail: dyliu@sklec.ecnu.edu.cn

on the biases in TN and  $\delta^{15}\text{N}$  values after acidification, and it is necessary to evaluate the biases according to the  $\text{CaCO}_3$  contents of the sediments.

Global mean  $\text{CaCO}_3$  content in marine surface sediments is approximately 34.8%, with a wide range of approximately 0%–80% (Ridgwell and Hargreaves, 2007). For examples, the samples from north and west Pacific Ocean contain approximately 0%–20%  $\text{CaCO}_3$ , but  $\text{CaCO}_3$  contents in the samples from Atlantic Ocean can reach to 80% (Ridgwell and Hargreaves, 2007). In this study, TN and  $\delta^{15}\text{N}$  values in 236 marine sediment samples with higher  $\text{CaCO}_3$  contents (20%–40%) from the coastal waters in the western Australia and 273 samples with lower  $\text{CaCO}_3$  contents (0%–16%) from the coastal waters in China's seas were measured, respectively, under acidified and non-acidified conditions. The biases of TN and  $\delta^{15}\text{N}$  values were evaluated and the correlation between the biases and  $\text{CaCO}_3$  contents in the

sediments were examined to understand the impact of  $\text{CaCO}_3$  contents.

## 2 Materials and methods

### 2.1 Sample collection

Two sediment cores were collected from the coast of King George River (KGR) in the northwestern Australia, and 236 sediment samples were obtained by slicing at 1-cm interval. Two and one sediment cores were collected from the Bohai Sea and Yellow Sea in China, respectively, and 221 sediment samples were obtained by slicing at 1-cm interval. In addition, 52 surface marine sediment samples were collected from the South Yellow Sea, China (Table 1). Totally, 509 marine sediment samples were stored in a freezer at  $-20^\circ\text{C}$  preparing for the analysis of TOC, TN,  $\delta^{15}\text{N}$  values and  $\text{CaCO}_3$  contents.

**Table 1.** Description of sample property

Samples identifier	Location	Number of samples	Description	TOC <sup>1)</sup> /%	TN <sup>2)</sup> %	C/N <sup>2)</sup>	$\delta^{15}\text{N}^{2)/\text{‰}}$	$\text{CaCO}_3/\%$
KGR6	13°55'13.44"S, 127°19'13.8"E	114	Down-core marine sediment from King George River, northwestern Australia (11.8 m depth)	0.40–0.75	0.03–0.11	8–19	3.6–5.7	20–31
KGR10	13°53'08.88"S, 127°08'22.56"E	122	Down-core marine sediment from King George River, northwestern Australia (11.2 m depth)	0.38–0.83	0.04–0.12	7–22	4.0–7.8	23–40
B62-2	38°52'12"N, 119°40'12"E	76	Down-core marine sediment from Bohai Sea, China (24 m depth)	0.46–0.92	0.06–0.12	6–12	4.9–6.4	1–7
B24-1	38°09'27.6"N, 122°29'0.24"E	63	Down-core marine sediment from North Yellow Sea, China (45 m depth)	0.68–1.14	0.09–0.18	7–12	4.5–5.6	1–5
H10-1	34°59'24"N, 123°29'56.04"E	82	Down-core marine sediment from South Yellow Sea, China (78 m depth)	0.20–1.09	0.03–0.14	7–14	3.8–6.2	1–7
SMS	–	52	Surface marine sediment from South Yellow Sea, China	0.14–1.18	0.02–0.19	6–21	3.7–6.0	1–16

Note: <sup>1)</sup> Measurement on acidified sample; <sup>2)</sup> measurement on unacidified whole sample.

### 2.2 Analytical procedures

TOC, TN and  $\delta^{15}\text{N}$  values were measured using a continuous-flow isotope-ratio mass spectrometer (Delta V Advantage, Thermo Scientific, Germany) coupled to an elemental analyzer (Flash EA 1112, Thermo Scientific, Italy) in Littoral Environment et Sociétés (LIENSs)-UMR7266, France. Firstly, all samples were freeze-dried and homogenized by grinding. Each sample was divided into two shares, using for the pretreatment with and without acidification, respectively. One share was acidified by adding 1 mL of 0.5 mol/L HCl (or 1 mol/L HCl) to every 100 mg of samples. The acidified samples were dried at  $<60^\circ\text{C}$ , then mixed with 1 mL Milli-Q waters and freeze-dried again. The freeze-dried samples were weighed into 5 mm×8 mm tin capsules, and the prepared capsules were used for the measurement of total organic carbon (TOC), total nitrogen ( $\text{TN}_{\text{acidified}}$ ) and stable nitrogen isotope ( $\delta^{15}\text{N}_{\text{acidified}}$ ). The corrections for TOC and  $\text{TN}_{\text{acidified}}$  values were performed because the weight of samples had increased after decarbonation (formation of  $\text{CaCl}_2$ ). The other sample share, using for non-acidification treatment, was weighed directly and placed into 5 mm×8 mm tin capsules for the measurement of total carbon (TC), total nitrogen ( $\text{TN}_{\text{unacidified}}$ ) and stable nitrogen isotope ( $\delta^{15}\text{N}_{\text{unacidified}}$ ). Based on the results of TOC and TC, the  $\text{CaCO}_3$  contents were estimated using  $\text{CaCO}_3 = (\text{TC} - \text{TOC}) \times 8.33$  (Stein et al., 1994; Schubert and Calvert, 2001).

The isotopic composition results are expressed relative to atmospheric  $\text{N}_2$  for  $\delta^{15}\text{N}$ . Replicate measurements of an acetanilide standard (Thermo Scientific) indicated that the  $\delta^{15}\text{N}$  analytical errors were less than 0.1‰. The total organic matter and total nitrogen weight data obtained as part of the stable isotope analyses. Replicate measurements of a certified reference materi-

al (Low Organic Content Soil, Elemental MicroAnalysis, UK) indicated relative errors of 0.025% and 0.002% for total organic carbon and total nitrogen, respectively.

### 2.3 Statistical analysis

We tested the differences of TN and  $\delta^{15}\text{N}$  values between non-acidified and acidified samples, using software SPSS 16.0 (*t*-test). The correlation between these differences and  $\text{CaCO}_3$  contents were analyzed respectively using SigmaPlot 12.0. The biases in TN and  $\delta^{15}\text{N}$  values after acidification were calculated, respectively, using the following equations:

$$\Delta\text{N (of the TN)} = \frac{\text{TN}_{\text{unacidified}} - \text{TN}_{\text{acidified}}}{\text{TN}_{\text{unacidified}}} \times 100, \quad (1)$$

$$\Delta^{15}\text{N} = \delta^{15}\text{N}_{\text{unacidified}} - \delta^{15}\text{N}_{\text{acidified}}. \quad (2)$$

## 3 Results and discussion


### 3.1 $\text{CaCO}_3$ contents of all samples

The  $\text{CaCO}_3$  contents were approximately 20%–40% and 1%–16% for the sediments collected from the northwestern Australia and China's seas, respectively (Table 1). Based on the  $\text{CaCO}_3$  contents of global marine sediments (Ridgwell and Hargreaves, 2007), the samples were divided into  $\text{CaCO}_3$ -poor samples ( $\text{CaCO}_3$  contents  $<20\%$ ) and  $\text{CaCO}_3$ -rich samples ( $\text{CaCO}_3$  contents  $>20\%$ ), respectively, for a comparison.


### 3.2 Effects of acid treatment on TN

The results of paired-sample *t*-test indicated a significant difference in TN contents ( $P=0.000<0.01$ , 2-tailed) between the samples measured with and without acid treatment. For the non-acidified samples, TN contents ranged from 0.02%–0.19%, while this range reduced to 0.01%–0.16% for the acidified samples. About 99% non-acidified vs. acidified TN values are located below the ideal 1:1 relation, indicating that acidification resulted in the losses of nitrogen contents from the samples (Fig. 1a). In ad-

dition, losses of nitrogen contents were on average higher in  $\text{CaCO}_3$ -rich samples compared to  $\text{CaCO}_3$ -poor samples after acidification. About 0%–40% and 10%–60% of nitrogen contents were lost from non-acidified  $\text{CaCO}_3$ -poor samples and  $\text{CaCO}_3$ -rich samples, respectively (Fig. 2a). Moreover, our data suggest a statistically significant positive correlation between the losses of nitrogen contents and the  $\text{CaCO}_3$  contents (Fig. 2a). These findings partially agree with Lohse et al. (2000) and Fernandes and Krull (2008), who found a particularly high losses of nitrogen



**Fig. 1.** Scatter plots of unacidified vs. acidified values for TN and  $\delta^{15}\text{N}$ . The ideal relation (1:1) is described by the solid line. The dashed lines indicate the statistic relation by regression analysis.



**Fig. 2.** Correlation between  $\text{CaCO}_3$  contents and the biases in TN and  $\delta^{15}\text{N}$  values after acidification.

**Table 2.** Comparison of differences caused by acidification in sedimentary TN and  $\delta^{15}\text{N}$  values with references

Sample type	$\text{CaCO}_3/\%$	$\Delta\text{N}/\%$	$\Delta^{15}\text{N}/\text{‰}$	Acidification method	Possible derives	Analysis mode	Reference
DSS <sup>1)</sup>	55–61	28–47	–	<i>in situ</i> $\text{H}_2\text{SO}_3$	$\text{H}_2\text{SO}_3$ volatilises nitrogenous organic matter	single <sup>6)</sup>	Lohse et al. (2000)
SMS <sup>2)</sup>	–	11–29	–	direct HCl acidification	nitrogen appears to be liberated during direct acidification	single	Ryba and Burgess (2002)
MPOM <sup>3)</sup>	<5	0	0	HCl fumes for 4 h	HCl fume treatment did not alter $\delta^{15}\text{N}$ and N values for low carbonate contents	dual <sup>7)</sup>	Lorrain et al. (2003)
DSS <sup>1)</sup>	30	0.3	0.005	<i>in situ</i> 2 mol/L HCl	weak HCl (2 mol/L) did not significantly affect $\delta^{15}\text{N}$ and TN values	dual	Kennedy et al. (2005)
SMS <sup>2)</sup>	>30	30–64	1–2	HCl washing	losses through volatilisation or solubilisation (acid-soluble compounds)	single	Fernandes and Krull (2008)
SES <sup>4)</sup>	<10	19–40	0–1				
SOILB <sup>5)</sup>	–	15	0.7	<i>in situ</i> 5% HCl	loss of soluble organic and/or inorganic N species, e.g., $^{15}\text{N}$ -rich species (e.g., protein, ammonium)	single	Brodie et al. (2011a, b)

Note: Single or dual indicate separate or simultaneous C and N analyses; <sup>1)</sup> deep-sea sediment; <sup>2)</sup> surface marine sediment; <sup>3)</sup> marine suspended particulate organic matter; <sup>4)</sup> surface estuarine sediment; <sup>5)</sup> international soil standard; <sup>6)</sup> measurement on unacidified whole sample for TN and  $\delta^{15}\text{N}$ ; <sup>7)</sup> measurement on acidified sample for TN and  $\delta^{15}\text{N}$ .

contents in  $\text{CaCO}_3$ -rich (>30%) samples, up to 64% after acid treatment (Table 2), but no significant correlation between the loss of nitrogen contents and the  $\text{CaCO}_3$  contents.

The loss of nitrogen contents from acidified samples was observed in the present and previous studies, although the acidification methods were different (Table 2). In our study, we used an acid treatment with 0.5 mol/L or 1 mol/L HCl in glass tubes (Peng et al., 2015), which avoided the loss of acid-soluble organic matter and/or fine particles by rinsing process (rinse method) and intensive effervescence (*in situ* acidification). This indicates that volatilisation of nitrogen compounds seems to be the most reasonable explanation for our results, and further investigation is still need for the explanation of the mechanism. Previous studies have demonstrated that in the  $\text{CaCO}_3$ -rich samples the fraction of proteinaceous compounds absorbed on carbonate surface layers became unstable with the increasing of acid addition, and this acid-soluble component was lost during carbonate dissolution (Fernandes and Krull, 2008). Meanwhile, this fraction of acid-soluble organic matter is greater in more calcareous sediments (Froelich, 1980). This may explain the higher losses observed in the  $\text{CaCO}_3$ -rich samples compared to the  $\text{CaCO}_3$ -poor samples.

### 3.3 Effects of acid treatment on $\delta^{15}\text{N}$

Nitrogen isotope ( $\delta^{15}\text{N}$ ) values of samples measured with and without acidification are shown in Fig. 1b. There have significant shifts in  $\delta^{15}\text{N}$  values between acidified and non-acidified samples, and these shifts have statistically significant positive correlation with the  $\text{CaCO}_3$  contents, indicating that losses of acid-soluble nitrogen compounds with different  $\delta^{15}\text{N}$  values (Fig. 2b). For  $\text{CaCO}_3$ -poor sediments the measured  $\delta^{15}\text{N}$  values after acid treatment varied in the range of  $-1\text{‰}$  to  $6\text{‰}$  (Fig. 1b). The shift of  $\delta^{15}\text{N}$  values of these samples after acidification (up to  $3\text{‰}$ ) is similar to the deviation ( $0\text{‰}$ – $2\text{‰}$ ) observed by Fernandes and Krull (2008) (Table 2). For  $\text{CaCO}_3$ -rich samples about 40% of the measured  $\delta^{15}\text{N}$  values were significant negative after acidification (down to  $-9\text{‰}$ , Fig. 1b), and bias in  $\delta^{15}\text{N}$  values (up to  $14\text{‰}$ ) were remarkable compared to non-acidified samples (Fig. 2b). Schmidt and Gleixner (2005) also found some samples with unusual negative  $\delta^{15}\text{N}$  values (e.g.,  $-4.24\text{‰}$  for a podzol Sample 6) after acid treatment due to the accumulation of coal-derived products. However, it is really difficult to understand what cause such depletion for part of  $\text{CaCO}_3$ -rich samples, since there is no rinse of the samples, which could trigger a loss of  $^{15}\text{N}$ -enriched compounds. Volatilisation of nitrogenous compounds like chloramine should lead to an enrichment of  $\delta^{15}\text{N}$  values not depletion.

In addition, our data suggest that isotopic shifts were not consistent for all samples (Fig. 2b). The depletion of  $\delta^{15}\text{N}$  values was observed in 95% samples after acid treatment reflecting the loss of  $^{15}\text{N}$ -enriched compounds such as protein and ammonium (Brodie et al., 2011a, b) (Fig. 2b). The loss of heavy isotope during acid treatment had previously been reported by other studies (Table 2). However, about 5% acidified samples (originating in  $\text{CaCO}_3$ -poor sediments collected from the Bohai Sea) detected a  $0\text{‰}$ – $1\text{‰}$  increase signature in  $\delta^{15}\text{N}$  values, suggesting the loss of  $^{14}\text{N}$ -enriched compounds. Harris et al. (2001) observed a small increase ( $0.04\text{‰}$ – $0.11\text{‰}$ ) in  $\delta^{15}\text{N}$  values of acidified soil samples following HCl fumigation. This contrasting shifts (depletion or enrichment) in  $\delta^{15}\text{N}$  values probably influenced by the organic matter composition and clay mineralogy of samples (Baldock and Skjemstad, 2000; Schubert and Calvert, 2001; Fernandes and Krull, 2008), but more work need to do to identify differences in acid-soluble compounds, inorganic nitrogen and mineralogy of

samples containing different  $\text{CaCO}_3$  contents during acid treatment.

### 4 Conclusions

Our results suggest that the analysis of nitrogen contents and its isotopic composition should be measured on non-acidified whole sample, because acidification to remove carbonates in marine sediments leads to significant nitrogen losses through volatilisation and isotopic shifts, particularly for samples with high  $\text{CaCO}_3$  contents (>20%). Meanwhile, there was a statistically significant positive correlation between the  $\text{CaCO}_3$  contents and the biases in nitrogen contents and its isotopic composition after acid treatment (Fig. 2), indicating that the  $\text{CaCO}_3$  contents in the sample play an important role in determining what extent the nitrogen fraction may have been affected by acid treatment.

### References

- Baldock J A, Skjemstad J O. 2000. Role of the soil matrix and minerals in protecting natural organic materials against biological attack. *Organic Geochemistry*, 31(7–8): 697–710
- Brodie C R, Heaton T H E, Leng M J, et al. 2011a. Evidence for bias in measured  $\delta^{15}\text{N}$  values of terrestrial and aquatic organic materials due to pre-analysis acid treatment methods. *Rapid Communications in Mass Spectrometry*, 25(8): 1089–1099
- Brodie C R, Leng M J, Casford J S L, et al. 2011b. Evidence for bias in C and N concentrations and  $\delta^{13}\text{C}$  composition of terrestrial and aquatic organic materials due to pre-analysis acid preparation methods. *Chemical Geology*, 282(3–4): 67–83
- Bunn S E, Loneragan N R, Kempster M A. 1995. Effects of acid washing on stable isotope ratios of C and N in penaeid shrimp and seagrass: Implications for food-web studies using multiple stable isotopes. *Limnology and Oceanography*, 40(3): 622–625
- Fernandes M, Krull E. 2008. How does acid treatment to remove carbonates affect the isotopic and elemental composition of soils and sediments?. *Environmental Chemistry*, 5(1): 33–39
- Froelich P N. 1980. Analysis of organic carbon in marine sediment. *Limnology and Oceanography*, 25(3): 564–572
- Harris D, Horwath W R, Van Kessel C. 2001. Acid fumigation of soils to remove carbonates prior to total organic carbon or carbon-13 isotopic analysis. *Soil Science Society of America*, 65(6): 1853–1856
- Hu Jianfang, Peng Ping'an, Jia Guodong, et al. 2006. Distribution and sources of organic carbon, nitrogen and their isotopes in sediments of the subtropical Pearl River estuary and adjacent shelf, Southern China. *Marine Chemistry*, 98(2–4): 274–285
- Jacob U, Mintenbeck K, Brey T, et al. 2005. Stable isotope food web studies: a case for standardized sample treatment. *Marine Ecology Progress Series*, 287: 251–253
- Jaschinski S, Hansen T, Sommer U. 2008. Effects of acidification in multiple stable isotope analyses. *Limnology and Oceanography: Methods*, 6(1): 12–15
- Kennedy P, Kennedy H, Papadimitriou S. 2005. The effect of acidification on the determination of organic carbon, total nitrogen and their stable isotopic composition in algae and marine sediment. *Rapid Communications in Mass Spectrometry*, 19(8): 1063–1068
- Lohse L, Kloosterhuis R T, De Stigter H C, et al. 2000. Carbonate removal by acidification causes loss of nitrogenous compounds in continental margin sediments. *Marine Chemistry*, 69(3–4): 193–201
- Lorrain A, Savoye N, Chauvaud L, et al. 2003. Decarbonation and preservation method for the analysis of organic C and N contents and stable isotope ratios of low-carbonated suspended particulate material. *Analytica Chimica Acta*, 491(2): 125–133
- Middelburg J J, Nieuwenhuize J. 1998. Carbon and nitrogen stable isotopes in suspended matter and sediments from the Schelde Estuary. *Marine Chemistry*, 60(3–4): 217–225
- Peng Yajun, Wang Yujue, Liu Dongyan, et al. 2015. Acid treatment effects on the carbon stable isotope values of marine sediments.

- Acta Oceanologica Sinica* (in Chinese), 37(12): 77–84
- Ridgwell A, Hargreaves J C. 2007. Regulation of atmospheric CO<sub>2</sub> by deep-sea sediments in an Earth system model. *Global Biogeochemical Cycles*, 21(2): GB2008, doi: [10.1029/2006GB002764](https://doi.org/10.1029/2006GB002764)
- Ryba S A, Burgess R M. 2002. Effects of sample preparation on the measurement of organic carbon, hydrogen, nitrogen, sulfur, and oxygen concentrations in marine sediments. *Chemosphere*, 48(1): 139–147
- Savage C. 2005. Tracing the influence of sewage nitrogen in a coastal ecosystem using stable nitrogen isotopes. *AMBIO: A Journal of the Human Environment*, 34(2): 145–150
- Schlacher T A, Connolly R M. 2014. Effects of acid treatment on carbon and nitrogen stable isotope ratios in ecological samples: a review and synthesis. *Methods in Ecology and Evolution*, 5(6): 541–550
- Schmidt M W I, Gleixner G. 2005. Carbon and nitrogen isotope composition of bulk soils, particle-size fractions and organic material after treatment with hydrofluoric acid. *European Journal of Soil Science*, 56(3): 407–416
- Schubert C J, Calvert S E. 2001. Nitrogen and carbon isotopic composition of marine and terrestrial organic matter in Arctic Ocean sediments: implications for nutrient utilization and organic matter composition. *Deep Sea Research Part I: Oceanographic Research Papers*, 48(3): 789–810
- Schubert C J, Nielsen B. 2000. Effects of decarbonation treatments on  $\delta^{13}\text{C}$  values in marine sediments. *Marine Chemistry*, 72(1): 55–59
- Serrano O, Serrano L, Mateo M A, et al. 2008. Acid washing effect on elemental and isotopic composition of whole beach arthropods: implications for food web studies using stable isotopes. *Acta Oecologica*, 34(1): 89–96
- Stein R, Grobe H, Wahsner M. 1994. Organic carbon, carbonate, and clay mineral distributions in eastern central Arctic Ocean surface sediments. *Marine Geology*, 119(3–4): 269–285
- Sweeney R E, Kaplan I R. 1980. Natural abundances of  $^{15}\text{N}$  as a source indicator for near-shore marine sedimentary and dissolved nitrogen. *Marine Chemistry*, 9(2): 81–94
- Tucker J, Sheats N, Giblin A E, et al. 1999. Using stable isotopes to trace sewage-derived material through Boston harbor and Massachusetts bay. *Marine Environmental Research*, 48(4–5): 353–375
- Voss M, Larsen B, Leivuori M, et al. 2000. Stable isotope signals of eutrophication in Baltic Sea sediments. *Journal of Marine Systems*, 25(3–4): 287–298
- Wang Yujue, Liu Dongyan, Richard P, et al. 2013. A geochemical record of environmental changes in sediments from Sishili Bay, northern Yellow Sea, China: Anthropogenic influence on organic matter sources and composition over the last 100 years. *Marine Pollution Bulletin*, 77(1–2): 227–236