


Crystal structure of 2,6-bis[(imidazol-1-yl)methyl]-4-chlorophenol — terephthalic acid (1:1), $C_{14}H_{13}ClN_4O$ — $C_8H_6O_4$

Wenhai Wang and Lingxin Chen*

Key Laboratory of Coastal Environmental Processes, Yantai Institute of Coastal Zone Research, Chinese Academy of Sciences, Yantai 264003, P. R. China

Received April 25, 2011, accepted and available on-line August 31, 2011; CCDC no. 1267/3546



Abstract

$C_{22}H_{19}ClN_4O_5$, triclinic, $P\bar{1}$ (no. 2), $a = 7.207(2)$ Å, $b = 12.087(7)$ Å, $c = 12.213(4)$ Å, $\alpha = 103.01(4)$ °, $\beta = 90.08(3)$ °, $\gamma = 102.20(3)$ °, $V = 1011.8$ Å³, $Z = 2$, $R_{gt}(F) = 0.044$, $wR_{ref}(F^2) = 0.113$, $T = 290$ K.

Source of material

2,6-bis[(imidazol-1-yl)methyl]-4-chlorophenol (BICP) was prepared according to [1]. The mixture of BICP (0.05 mmol, 14.4 mg) and terephthalic acid (TPA, 0.05 mmol, 8.3 mg) was dissolved in methanol (15 mL) with stirring. Upon slow evaporation of solvent, colorless block single crystals suitable for X-ray analysis were obtained after two weeks (yield 68 %).

Experimental details

All H atoms were located in a difference Fourier map and refined with $U_{iso}(H) = 1.2 U_{eq}(C)$ and $1.5 U_{eq}(O)$.

Discussion

Intermolecular hydrogen bonds have proven to be an ideal and efficient tool in the design and construction of organic crystals because of their strength and directional properties [2]. The architectures assembled from organic ligands and various hydrogen bonding donors has been rapidly developed because of their fascinating structural diversity and potential applications as func-

tional materials [3,4]. Carboxylic acids which can form strong and directional hydrogen bonds are frequently chosen as building blocks for crystal engineering, and a variety of co-crystals have been synthesized by assembling carboxylic acids and organic ligands bearing *N*-donors, such as pyridine-based ligands [5–7]. Imidazoles, which are also *N*-donor compounds, are used to construct organic crystals, but only a few ones composed of carboxylic acids and diimidazole compounds have been reported in the literature to date [8–10].

The asymmetric unit of the title crystal structure contains one TPA and one BICP molecule. Two imidazole rings of the BICP moiety are located at the different sides of the benzene ring with the angles $N1-C7-C3 = 111.3$ ° and $N3-C11-C1 = 111.0$ °. The carboxyl group of TPA interacts with the imidazol-1-yl nitrogen atom of BICP via $O2-H2O$ — $N2$ and $O5-H5$ — $N2$ hydrogen bonds, and thus the hydrogen bonds further propagate the acid-base subunits into an infinite one-dimensional zigzag chain. Meanwhile, every two chains are combined into a double chain by $O1-H1$ — $O3$ hydrogen bonds between the hydroxyl group of BICP and the carboxyl group of TPA. In addition, the $C11$ and $C14$ of the BICP molecule and the $O3$ of the carboxylic group could form $C11-H11A$ — $O3$ and $C14-H14$ — $O3$ hydrogen bonds, which is another contributor for the combination of two single chains. Further analysis reveals that the weak intermolecular C—H π interaction also contributes to the crystal packing arrangement.

Table 1. Data collection and handling.

Crystal:	colourless block, size 0.18 0.22 0.26 mm
Wavelength:	Mo $K\alpha$ radiation (0.71073 Å)
μ :	2.34 cm ⁻¹
Diffractometer, scan mode:	CAD4, $\omega/2\theta$
$2\theta_{max}$:	50.08°
$N(hkl)_{measured}$, $N(hkl)_{unique}$:	3539, 3539
Criterion for I_{obs} , $N(hkl)_{gt}$:	$I_{obs} > 2 \sigma(I_{obs})$, 3539
$N(param)_{refined}$:	365
Programs:	SHELXS-97, SHELXL-97 [11]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	x	y	z	U_{iso}
H(4)	2 <i>i</i>	0.716(4)	1.149(3)	0.071(2)	0.047
H(6)	2 <i>i</i>	0.790(4)	0.949(2)	0.143(2)	0.051
H(7B)	2 <i>i</i>	0.630(4)	1.058(3)	0.269(2)	0.044
H(7A)	2 <i>i</i>	0.622(4)	0.926(2)	0.321(2)	0.039
H(8)	2 <i>i</i>	0.911(4)	1.209(2)	0.296(2)	0.047
H(9)	2 <i>i</i>	1.278(5)	1.015(3)	0.410(3)	0.060
H(10)	2 <i>i</i>	0.974(4)	0.883(2)	0.361(2)	0.046

* Correspondence author (e-mail: lxchen@yic.ac.cn)

Table 2. Continued.

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(11B)	<i>2i</i>	0.844(4)	0.758(2)	0.075(2)	0.038
H(11A)	<i>2i</i>	0.817(4)	0.694(2)	0.057(2)	0.035
H(12)	<i>2i</i>	0.555(5)	0.725(3)	0.193(3)	0.070
H(13)	<i>2i</i>	0.167(4)	0.528(3)	0.048(2)	0.054
H(14)	<i>2i</i>	0.484(5)	0.602(3)	0.138(3)	0.080
H(18)	<i>2i</i>	0.123(4)	0.343(2)	0.407(2)	0.041

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
C(1)	<i>2i</i>	0.7568(3)	0.8600(2)	0.0233(2)	0.025(1)	0.033(2)	0.035(1)	0.004(1)	0.007(1)	0.012(1)
C(2)	<i>2i</i>	0.7322(3)	0.8723(2)	0.1332(2)	0.026(1)	0.029(1)	0.032(1)	0.001(1)	0.005(1)	0.007(1)
C(3)	<i>2i</i>	0.7175(3)	0.9811(2)	0.1509(2)	0.024(1)	0.038(2)	0.034(1)	0.005(1)	0.005(1)	0.014(1)
C(4)	<i>2i</i>	0.7306(4)	1.0743(2)	0.0608(2)	0.030(2)	0.028(2)	0.047(2)	0.005(1)	0.006(1)	0.013(1)
C(5)	<i>2i</i>	0.7575(4)	1.0620(2)	0.0471(2)	0.029(2)	0.030(2)	0.043(2)	0.003(1)	0.006(1)	0.005(1)
C(6)	<i>2i</i>	0.7711(4)	0.9558(2)	0.0659(2)	0.028(2)	0.040(2)	0.032(2)	0.003(1)	0.004(1)	0.011(1)
Cl	<i>2i</i>	0.7776(1)	1.18079(7)	0.16060(6)	0.0610(5)	0.0395(4)	0.0522(5)	0.0103(4)	0.0009(4)	0.0061(3)
C(7)	<i>2i</i>	0.6968(4)	0.9968(3)	0.2694(2)	0.031(2)	0.044(2)	0.042(2)	0.004(1)	0.003(1)	0.022(1)
N(1)	<i>2i</i>	0.8827(3)	1.0328(2)	0.3143(2)	0.034(1)	0.037(1)	0.033(1)	0.007(1)	0.005(1)	0.016(1)
C(8)	<i>2i</i>	0.9717(4)	1.1423(3)	0.3157(2)	0.039(2)	0.041(2)	0.040(2)	0.008(1)	0.010(1)	0.021(1)
N(2)	<i>2i</i>	1.1430(3)	1.1458(2)	0.3539(2)	0.040(2)	0.049(2)	0.044(1)	0.005(1)	0.008(1)	0.023(1)
C(9)	<i>2i</i>	1.1644(5)	1.0335(3)	0.3777(2)	0.043(2)	0.058(2)	0.042(2)	0.018(2)	0.011(1)	0.022(2)
C(10)	<i>2i</i>	1.0064(4)	0.9628(3)	0.3537(2)	0.048(2)	0.039(2)	0.040(2)	0.011(2)	0.008(1)	0.014(1)
O(1)	<i>2i</i>	0.7169(3)	0.7858(2)	0.2270(2)	0.056(1)	0.032(1)	0.033(1)	0.012(1)	0.0043(9)	0.0075(9)
C(11)	<i>2i</i>	0.7648(4)	0.7453(2)	0.0033(2)	0.040(2)	0.037(2)	0.033(2)	0.008(1)	0.006(1)	0.013(1)
N(3)	<i>2i</i>	0.5724(3)	0.6802(2)	0.0195(2)	0.040(1)	0.030(1)	0.031(1)	0.005(1)	0.008(1)	0.013(1)
C(12)	<i>2i</i>	0.4860(5)	0.6840(3)	0.1174(3)	0.060(2)	0.040(2)	0.037(2)	0.012(2)	0.016(2)	0.019(1)
N(4)	<i>2i</i>	0.3101(4)	0.6223(2)	0.0991(2)	0.054(2)	0.043(2)	0.053(2)	0.008(1)	0.022(1)	0.026(1)
C(13)	<i>2i</i>	0.2848(5)	0.5764(3)	0.0144(3)	0.049(2)	0.047(2)	0.052(2)	0.001(2)	0.006(2)	0.017(2)
C(14)	<i>2i</i>	0.4452(4)	0.6119(3)	0.0639(3)	0.050(2)	0.048(2)	0.033(2)	0.002(2)	0.007(1)	0.013(1)
C(19)	<i>2i</i>	0.0516(4)	0.4280(2)	0.3465(2)	0.038(2)	0.025(1)	0.025(1)	0.001(1)	0.005(1)	0.001(1)
C(22)	<i>2i</i>	0.0906(4)	0.4752(2)	0.2698(2)	0.037(2)	0.028(2)	0.033(1)	0.000(1)	0.010(1)	0.001(1)
O(2)	<i>2i</i>	0.0597(3)	0.5624(2)	0.2347(2)	0.056(1)	0.041(1)	0.057(1)	0.014(1)	0.025(1)	0.027(1)
O(3)	<i>2i</i>	0.2287(3)	0.4313(2)	0.2438(2)	0.054(1)	0.043(1)	0.063(1)	0.019(1)	0.034(1)	0.022(1)
C(18)	<i>2i</i>	0.0037(4)	0.3559(2)	0.4104(2)	0.032(2)	0.035(2)	0.030(1)	0.007(1)	0.006(1)	0.005(1)
C(17)	<i>2i</i>	0.1342(4)	0.3070(2)	0.4771(2)	0.038(2)	0.037(2)	0.032(2)	0.008(1)	0.005(1)	0.015(1)
C(16)	<i>2i</i>	0.3173(4)	0.3270(2)	0.4803(2)	0.033(2)	0.034(2)	0.029(1)	0.002(1)	0.006(1)	0.008(1)
C(15)	<i>2i</i>	0.4551(4)	0.2690(2)	0.5516(2)	0.035(2)	0.045(2)	0.039(2)	0.006(1)	0.008(1)	0.017(1)
O(4)	<i>2i</i>	0.4117(3)	0.2098(2)	0.6107(2)	0.043(1)	0.112(2)	0.096(2)	0.021(1)	0.022(1)	0.080(2)
O(5)	<i>2i</i>	0.6258(3)	0.2881(2)	0.5475(2)	0.038(1)	0.070(2)	0.070(2)	0.017(1)	0.022(1)	0.044(1)
C(21)	<i>2i</i>	0.3660(4)	0.3994(2)	0.4168(2)	0.030(2)	0.036(2)	0.036(2)	0.007(1)	0.006(1)	0.009(1)
C(20)	<i>2i</i>	0.2316(4)	0.4506(2)	0.3515(2)	0.040(2)	0.032(2)	0.029(1)	0.005(1)	0.004(1)	0.010(1)

Acknowledgments. This work was financially supported by the National Natural Science Foundation of China (grant no. 20975089), the Department of Science and Technology of Shandong Province (grant no. 2008GG20005005) and the 100 Talents Program of the Chinese Academy of Sciences.

References

- Zhou, C. H.; Zhang, Z. J.; Yuan, D. Q.; Xie, R. G.; Zhao, H. M.: Synthesis of New Chiral Macroyclic Tetraoxo Polyamines Containing Pyridine Ring and Functional Arms. *Synth Commun.* **24** (1994) 47-52.
- Aakeröy, C. B.; Seddon, K. R.: The Hydrogen Bond and Crystal Engineering. *Chem. Soc. Rev.* **22** (1993) 397-407.
- MacGillivray, S. L. R.; Atwood, J. L.: A Chiral Spherical Molecular Assembly Held Together by 60 Hydrogen Bonds. *Nature* **389** (1997) 469-472.
- Corna, A.; Rey, F.; Rius, J.; Sabater, M. J.; Valencla, S.: Supramolecular Self-assembled Molecules as Organic Directing Agent for Synthesis of Zeolites. *Nature* **431** (2004) 287-290.
- Bhogala, B. R.; Ashwini N.: Cocrystals of 1,3,5-Cyclohexanetricarboxylic Acid with 4,4'-Bipyridine Homologues: Acid-Pyridine Hydrogen Bonding in Neutral and Ionic Complexes. *Cryst. Growth Des.* **3** (2003) 547-554.
- Childs, S. L.; Hardcastle, K. I.: Cocrystals of Piroxicam with Carboxylic. *Cryst. Growth Des.* **7** (2007) 1291-1304.
- Ramkinkar S.; Kumar, B.: Two-Dimensional Organic Brick-Wall Layers as Hosts for the Inclusion and Study of Aromatics Ensembles: Acid-Pyridine and Acid-Carbonyl Synthons for Multicomponent Materials. *Cryst. Growth Des.* **9** (2009) 4969-4978.
- MacDonald, J. C.; Dorrestein, P. C.; Pilley, M. M.: Design of Supramolecular Layers via Self-Assembly of Imidazole and Carboxylic Acids. *Cryst. Growth Des.* **1** (2001) 29-38.
- Cheruzel, L. E.; Mashuta, M. S.; Buchanan, R. M.: A Supramolecular Assembly of Side-by-side Polyimidazole Tripod Coils Stabilized by π-π Stacking and Unique Boric Acid Templated Hydrogen Bonding Interactions. *Chem. Commun.* (2005) 2223-2225.
- Aakeröy, C. B.; Salmon, D. J.; Smith, M. M.; Desper, J.: Cyanophenylloximes: Reliable and Versatile Tools for Hydrogen-Bond Directed Supramolecular Synthesis of Cocrystals. *Cryst. Growth. Des.* **6** (2006) 1033-1042.
- Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A64** (2008) 112-122.